

# Descriptive Statistics: Part 2/2 (Ch 3)

Will Landau

Iowa State University

January 24, 2013

# Outline

Descriptive  
Statistics: Part  
2/2 (Ch 3)

Will Landau

Boxplots

Boxplots

Quantile-Quantile (QQ) Plots

Quantile-Quantile  
(QQ) Plots

Theoretical Quantile-Quantile Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical Summaries

Numerical  
Summaries

Parameters

Parameters

# Generic Boxplot


## Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters



# Example: bullet data

Quantiles of the Bullet Penetration Depth Distributions

$i$	$\frac{i-.5}{20}$	$i$ th Smallest 230 Grain Data Point = $Q(\frac{i-.5}{20})$	$i$ th Smallest 200 Grain Data Point = $Q(\frac{i-.5}{20})$
1	.025	27.75	58.00
2	.075	37.35	58.65
3	.125	38.35	59.10
4	.175	38.35	59.50
5	.225	38.75	59.80
6	.275	39.75	60.70
7	.325	40.50	61.30
8	.375	41.00	61.50
9	.425	41.15	62.30
10	.475	42.55	62.65
11	.525	42.90	62.95
12	.575	43.60	63.30
13	.625	43.85	63.55
14	.675	47.30	63.80
15	.725	47.90	64.05
16	.775	48.15	64.65
17	.825	49.85	65.00
18	.875	51.25	67.75
19	.925	51.60	70.40
20	.975	56.00	71.70

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

## Example: bullet data (230-grain bullets)

$$Q(.25) = .5Q(.225) + .5Q(.275) = .5(38.75) + .5(39.75) = 39.25 \text{ mm}$$

$$Q(.5) = .5Q(.475) + .5Q(.525) = .5(42.55) + .5(42.90) = 42.725 \text{ mm}$$

$$Q(.75) = .5Q(.725) + .5Q(.775) = .5(47.90) + .5(48.15) = 48.025 \text{ mm}$$

So


$$IQR = 48.025 - 39.25 = 8.775 \text{ mm}$$

$$1.5IQR = 13.163 \text{ mm}$$

$$Q(.75) + 1.5IQR = 61.188 \text{ mm}$$

$$Q(.25) - 1.5IQR = 26.087 \text{ mm}$$

# Example: bullet data



Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

# Outline

Descriptive  
Statistics: Part  
2/2 (Ch 3)

Will Landau

Boxplots

Boxplots

Quantile-Quantile  
(QQ) Plots

Quantile-Quantile (QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Theoretical Quantile-Quantile Plots

Numerical  
Summaries

Numerical Summaries

Parameters

Parameters

- ▶ **Quantile-quantile (QQ) plot:** a scatterplot of the sorted values of one dataset on the sorted values of another dataset.
  - ▶ This plot is used to tell if the distributional shapes of the datasets are the same or different.
    - ▶ If the points in the plot lie in a straight line, the distributional shapes are the same.
    - ▶ Otherwise, the shapes are different.
  - ▶ The datasets must be univariate, numerical, and of the same size.

- ▶ **Quantile-quantile (QQ) plot:** a scatterplot of the sorted values of one dataset on the sorted values of another dataset.
  - ▶ This plot is used to tell if the distributional shapes of the datasets are the same or different.
    - ▶ If the points in the plot lie in a straight line, the distributional shapes are the same.
    - ▶ Otherwise, the shapes are different.
  - ▶ The datasets must be univariate, numerical, and of the same size.

- ▶ **Quantile-quantile (QQ) plot:** a scatterplot of the sorted values of one dataset on the sorted values of another dataset.
  - ▶ This plot is used to tell if the distributional shapes of the datasets are the same or different.
    - ▶ If the points in the plot lie in a straight line, the distributional shapes are the same.
    - ▶ Otherwise, the shapes are different.
  - ▶ The datasets must be univariate, numerical, and of the same size.

- ▶ **Quantile-quantile (QQ) plot:** a scatterplot of the sorted values of one dataset on the sorted values of another dataset.
  - ▶ This plot is used to tell if the distributional shapes of the datasets are the same or different.
    - ▶ If the points in the plot lie in a straight line, the distributional shapes are the same.
    - ▶ Otherwise, the shapes are different.
  - ▶ The datasets must be univariate, numerical, and of the same size.

- ▶ **Quantile-quantile (QQ) plot:** a scatterplot of the sorted values of one dataset on the sorted values of another dataset.
  - ▶ This plot is used to tell if the distributional shapes of the datasets are the same or different.
    - ▶ If the points in the plot lie in a straight line, the distributional shapes are the same.
    - ▶ Otherwise, the shapes are different.
  - ▶ The datasets must be univariate, numerical, and of the same size.

# Example: bullet data

Quantiles of the Bullet Penetration Depth Distributions

$i$	$\frac{i-.5}{20}$	$i$ th Smallest 230 Grain Data Point = $Q(\frac{i-.5}{20})$	$i$ th Smallest 200 Grain Data Point = $Q(\frac{i-.5}{20})$
1	.025	27.75	58.00
2	.075	37.35	58.65
3	.125	38.35	59.10
4	.175	38.35	59.50
5	.225	38.75	59.80
6	.275	39.75	60.70
7	.325	40.50	61.30
8	.375	41.00	61.50
9	.425	41.15	62.30
10	.475	42.55	62.65
11	.525	42.90	62.95
12	.575	43.60	63.30
13	.625	43.85	63.55
14	.675	47.30	63.80
15	.725	47.90	64.05
16	.775	48.15	64.65
17	.825	49.85	65.00
18	.875	51.25	67.75
19	.925	51.60	70.40
20	.975	56.00	71.70

Boxplots

Quantile-Quantile  
(QQ) Plots


Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters


## Example: bullet data

- ▶ I can make a QQ plot of the bullet data by plotting the sorted 200-grain depths against the sorted 230-grain depths.
- ▶ The points lie in approximately a straight line, so the 200-grain depths are similarly shaped in distribution to the 230-grain depths.



## Example: bullet data

- ▶ I can make a QQ plot of the bullet data by plotting the sorted 200-grain depths against the sorted 230-grain depths.
- ▶ The points lie in approximately a straight line, so the 200-grain depths are similarly shaped in distribution to the 230-grain depths.



Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

# Outline

Descriptive  
Statistics: Part  
2/2 (Ch 3)

Will Landau

Boxplots

Boxplots

Quantile-Quantile  
(QQ) Plots

Quantile-Quantile (QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Theoretical Quantile-Quantile Plots

Numerical  
Summaries

Numerical Summaries

Parameters

Parameters

# Theoretical quantile-quantile (QQ) plots

- ▶ **Theoretical quantile-quantile (QQ) plot:** a scatterplot with:

- ▶ The sorted values  $x_1, x_2, \dots, x_n$  of some real data set on the  $x$  axis.
- ▶  $Q(\frac{1-.5}{n}), Q(\frac{2-.5}{n}), \dots, Q(\frac{n-.5}{n})$  on the  $y$  axis.
  - ▶  $Q$  is some **theoretical quantile function**: the quantile function we would expect from a dataset if that dataset had a certain shape.

- ▶ Example theoretical quantile functions:

- ▶ "Standard" bell-shaped data should have:

$$Q(p) \approx 4.9(p^{0.14} - (1-p)^{0.14})$$

- ▶ "Exponentially distributed" data (a kind of highly right-skewed data) should have:

$$Q(p) \approx -\lambda^{-1} \log(1-p)$$

where  $\lambda$  is some constant.

# Theoretical quantile-quantile (QQ) plots

- ▶ **Theoretical quantile-quantile (QQ) plot:** a scatterplot with:

- ▶ The sorted values  $x_1, x_2, \dots, x_n$  of some real data set on the  $x$  axis.

- ▶  $Q(\frac{1-.5}{n}), Q(\frac{2-.5}{n}), \dots, Q(\frac{n-.5}{n})$  on the  $y$  axis.

- ▶  $Q$  is some **theoretical quantile function**: the quantile function we would expect from a dataset if that dataset had a certain shape.

- ▶ Example theoretical quantile functions:

- ▶ "Standard" bell-shaped data should have:

$$Q(p) \approx 4.9(p^{0.14} - (1-p)^{0.14})$$

- ▶ "Exponentially distributed" data (a kind of highly right-skewed data) should have:

$$Q(p) \approx -\lambda^{-1} \log(1-p)$$

where  $\lambda$  is some constant.

# Theoretical quantile-quantile (QQ) plots

- ▶ **Theoretical quantile-quantile (QQ) plot:** a scatterplot with:

- ▶ The sorted values  $x_1, x_2, \dots, x_n$  of some real data set on the  $x$  axis.
- ▶  $Q(\frac{1-.5}{n}), Q(\frac{2-.5}{n}), \dots, Q(\frac{n-.5}{n})$  on the  $y$  axis.
  - ▶  $Q$  is some **theoretical quantile function**: the quantile function we would expect from a dataset if that dataset had a certain shape.

- ▶ Example theoretical quantile functions:

- ▶ "Standard" bell-shaped data should have:

$$Q(p) \approx 4.9(p^{0.14} - (1-p)^{0.14})$$

- ▶ "Exponentially distributed" data (a kind of highly right-skewed data) should have:

$$Q(p) \approx -\lambda^{-1} \log(1-p)$$

where  $\lambda$  is some constant.

# Theoretical quantile-quantile (QQ) plots

- ▶ **Theoretical quantile-quantile (QQ) plot:** a scatterplot with:

- ▶ The sorted values  $x_1, x_2, \dots, x_n$  of some real data set on the  $x$  axis.
- ▶  $Q(\frac{1-.5}{n}), Q(\frac{2-.5}{n}), \dots, Q(\frac{n-.5}{n})$  on the  $y$  axis.
  - ▶  $Q$  is some **theoretical quantile function**: the quantile function we would *expect* from a dataset if that dataset had a certain shape.

- ▶ Example theoretical quantile functions:

- ▶ “Standard” bell-shaped data should have:

$$Q(p) \approx 4.9(p^{0.14} - (1-p)^{0.14})$$

- ▶ “Exponentially distributed” data (a kind of highly right-skewed data) should have:

$$Q(p) \approx -\lambda^{-1} \log(1-p)$$

where  $\lambda$  is some constant.

# Theoretical quantile-quantile (QQ) plots

- ▶ **Theoretical quantile-quantile (QQ) plot:** a scatterplot with:
  - ▶ The sorted values  $x_1, x_2, \dots, x_n$  of some real data set on the  $x$  axis.
  - ▶  $Q(\frac{1-.5}{n}), Q(\frac{2-.5}{n}), \dots, Q(\frac{n-.5}{n})$  on the  $y$  axis.
    - ▶  $Q$  is some **theoretical quantile function**: the quantile function we would *expect* from a dataset if that dataset had a certain shape.
- ▶ Example theoretical quantile functions:
  - ▶ “Standard” bell-shaped data should have:

$$Q(p) \approx 4.9(p^{0.14} - (1-p)^{0.14})$$

- ▶ “Exponentially distributed” data (a kind of highly right-skewed data) should have:

$$Q(p) \approx -\lambda^{-1} \log(1-p)$$

where  $\lambda$  is some constant.

# Theoretical quantile-quantile (QQ) plots

- ▶ **Theoretical quantile-quantile (QQ) plot:** a scatterplot with:

- ▶ The sorted values  $x_1, x_2, \dots, x_n$  of some real data set on the  $x$  axis.
- ▶  $Q(\frac{1-.5}{n}), Q(\frac{2-.5}{n}), \dots, Q(\frac{n-.5}{n})$  on the  $y$  axis.
  - ▶  $Q$  is some **theoretical quantile function**: the quantile function we would *expect* from a dataset if that dataset had a certain shape.

- ▶ Example theoretical quantile functions:

- ▶ “Standard” bell-shaped data should have:

$$Q(p) \approx 4.9(p^{0.14} - (1-p)^{0.14})$$

- ▶ “Exponentially distributed” data (a kind of highly right-skewed data) should have:

$$Q(p) \approx -\lambda^{-1} \log(1-p)$$

where  $\lambda$  is some constant.

# Theoretical quantile-quantile (QQ) plots

- ▶ **Theoretical quantile-quantile (QQ) plot:** a scatterplot with:

- ▶ The sorted values  $x_1, x_2, \dots, x_n$  of some real data set on the  $x$  axis.
- ▶  $Q(\frac{1-.5}{n}), Q(\frac{2-.5}{n}), \dots, Q(\frac{n-.5}{n})$  on the  $y$  axis.
  - ▶  $Q$  is some **theoretical quantile function**: the quantile function we would *expect* from a dataset if that dataset had a certain shape.

- ▶ Example theoretical quantile functions:

- ▶ “Standard” bell-shaped data should have:

$$Q(p) \approx 4.9(p^{0.14} - (1-p)^{0.14})$$

- ▶ “Exponentially distributed” data (a kind of highly right-skewed data) should have:

$$Q(p) \approx -\lambda^{-1} \log(1-p)$$

where  $\lambda$  is some constant.

# Normal quantile-quantile (QQ) Plots

- ▶ **Normal quantile-quantile (QQ) plot:** a theoretical QQ plot where the quantile function,  $Q$ , is the quantile function for “standard” bell-shaped (normally-distributed) data.
- ▶ If the points in a normal QQ plot are in a straight line, the dataset in question is bell-shaped. Otherwise, the data is not bell-shaped.

# Normal quantile-quantile (QQ) Plots

- ▶ **Normal quantile-quantile (QQ) plot:** a theoretical QQ plot where the quantile function,  $Q$ , is the quantile function for “standard” bell-shaped (normally-distributed) data.
- ▶ If the points in a normal QQ plot are in a straight line, the dataset in question is bell-shaped. Otherwise, the data is not bell-shaped.

# Example: towel breaking strength data

Descriptive  
Statistics: Part  
2/2 (Ch 3)

Will Landau

Breaking Strength and Standard Normal Quantiles

$i$	$\frac{i-.5}{10}$	$\frac{i-.5}{10}$ Breaking Strength Quantile	$\frac{i-.5}{10}$ Standard Normal Quantile
1	.05	7,583	-1.65
2	.15	8,527	-1.04
3	.25	8,572	-.67
4	.35	8,577	-.39
5	.45	9,011	-.13
6	.55	9,165	.13
7	.65	9,471	.39
8	.75	9,614	.67
9	.85	9,614	1.04
10	.95	10,688	1.65

Boxplots


Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries


Parameters

## Example: towel breaking strength data



- The points are roughly straight-line-shaped, so the breaking strength data is roughly bell-shaped.

## Normal QQ plot: 200-grain bullet penetration



- ▶ Since the points in the normal QQ plot are not quite arranged in a straight line, the 200-grain penetration depths are not quite bell-shaped. However, the departure from normality is not severe.
- ▶ The QQ plot of the bullet data from before revealed that the 200-grain depths had the same distributional shape as the 200-grain bullet depths. Thus, the 230-grain bullet data is not quite bell-shaped either.

- ▶ Since the points in the normal QQ plot are not quite arranged in a straight line, the 200-grain penetration depths are not quite bell-shaped. However, the departure from normality is not severe.
- ▶ The QQ plot of the bullet data from before revealed that the 200-grain depths had the same distributional shape as the 200-grain bullet depths. Thus, the 230-grain bullet data is not quite bell-shaped either.

# Outline

Descriptive  
Statistics: Part  
2/2 (Ch 3)

Will Landau

Boxplots

Boxplots

Quantile-Quantile  
(QQ) Plots

Quantile-Quantile (QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Theoretical Quantile-Quantile Plots

Numerical  
Summaries

Numerical Summaries

Parameters

Parameters

# Numerical summaries

## ▶ Numerical summary (statistic)

- ▶ A number or list of numbers calculated using the data (and only the data).
- ▶ Numerical summaries highlight important features of the data (shape, center, spread, outliers).

## ▶ Examples:

- ▶ Measures of center:
  - ▶ Arithmetic mean
  - ▶ Median
  - ▶ Mode
- ▶ Measures of spread:
  - ▶ Sample variance
  - ▶ Sample standard deviation
  - ▶ Range
  - ▶ IQR
- ▶ Measures of shape:
  - ▶ All the quantiles together
  - ▶ Skew (beyond the scope of the class)
  - ▶ Kurtosis (beyond the scope of the class)

# Numerical summaries

## ▶ Numerical summary (statistic)

- ▶ A number or list of numbers calculated using the data (and only the data).
- ▶ Numerical summaries highlight important features of the data (shape, center, spread, outliers).

## ▶ Examples:

### ▶ Measures of center:

- ▶ Arithmetic mean
- ▶ Median
- ▶ Mode

### ▶ Measures of spread:

- ▶ Sample variance
- ▶ Sample standard deviation
- ▶ Range
- ▶ IQR

### ▶ Measures of shape:

- ▶ All the quantiles together
- ▶ Skew (beyond the scope of the class)
- ▶ Kurtosis (beyond the scope of the class)

# Numerical summaries

## ▶ Numerical summary (statistic)

- ▶ A number or list of numbers calculated using the data (and only the data).
- ▶ Numerical summaries highlight important features of the data (shape, center, spread, outliers).

## ▶ Examples:

### ▶ Measures of center:

- ▶ Arithmetic mean
- ▶ Median
- ▶ Mode

### ▶ Measures of spread:

- ▶ Sample variance
- ▶ Sample standard deviation
- ▶ Range
- ▶ IQR

### ▶ Measures of shape:

- ▶ All the quantiles together
- ▶ Skew (beyond the scope of the class)
- ▶ Kurtosis (beyond the scope of the class)

# Numerical summaries

## ▶ Numerical summary (statistic)

- ▶ A number or list of numbers calculated using the data (and only the data).
- ▶ Numerical summaries highlight important features of the data (shape, center, spread, outliers).

## ▶ Examples:

### ▶ Measures of center:

- ▶ Arithmetic mean
- ▶ Median
- ▶ Mode

### ▶ Measures of spread:

- ▶ Sample variance
- ▶ Sample standard deviation
- ▶ Range
- ▶ IQR

### ▶ Measures of shape:

- ▶ All the quantiles together
- ▶ Skew (beyond the scope of the class)
- ▶ Kurtosis (beyond the scope of the class)

# Numerical summaries

## ▶ Numerical summary (statistic)

- ▶ A number or list of numbers calculated using the data (and only the data).
- ▶ Numerical summaries highlight important features of the data (shape, center, spread, outliers).

## ▶ Examples:

### ▶ Measures of center:

- ▶ Arithmetic mean
- ▶ Median
- ▶ Mode

### ▶ Measures of spread:

- ▶ Sample variance
- ▶ Sample standard deviation
- ▶ Range
- ▶ IQR

### ▶ Measures of shape:

- ▶ All the quantiles together
- ▶ Skew (beyond the scope of the class)
- ▶ Kurtosis (beyond the scope of the class)

# Numerical summaries

## ▶ Numerical summary (statistic)

- ▶ A number or list of numbers calculated using the data (and only the data).
- ▶ Numerical summaries highlight important features of the data (shape, center, spread, outliers).

## ▶ Examples:

- ▶ Measures of center:
  - ▶ Arithmetic mean
  - ▶ Median
  - ▶ Mode
- ▶ Measures of spread:
  - ▶ Sample variance
  - ▶ Sample standard deviation
  - ▶ Range
  - ▶ IQR
- ▶ Measures of shape:
  - ▶ All the quantiles together
  - ▶ Skew (beyond the scope of the class)
  - ▶ Kurtosis (beyond the scope of the class)

# Numerical summaries

## ▶ Numerical summary (statistic)

- ▶ A number or list of numbers calculated using the data (and only the data).
- ▶ Numerical summaries highlight important features of the data (shape, center, spread, outliers).

## ▶ Examples:

- ▶ Measures of center:
  - ▶ Arithmetic mean
  - ▶ Median
  - ▶ Mode
- ▶ Measures of spread:
  - ▶ Sample variance
  - ▶ Sample standard deviation
  - ▶ Range
  - ▶ IQR
- ▶ Measures of shape:
  - ▶ All the quantiles together
  - ▶ Skew (beyond the scope of the class)
  - ▶ Kurtosis (beyond the scope of the class)

# Numerical summaries

Descriptive  
Statistics: Part  
2/2 (Ch 3)

Will Landau

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

## ▶ Numerical summary (statistic)

- ▶ A number or list of numbers calculated using the data (and only the data).
- ▶ Numerical summaries highlight important features of the data (shape, center, spread, outliers).

## ▶ Examples:

- ▶ Measures of center:
  - ▶ Arithmetic mean
  - ▶ Median
  - ▶ Mode
- ▶ Measures of spread:
  - ▶ Sample variance
  - ▶ Sample standard deviation
  - ▶ Range
  - ▶ IQR
- ▶ Measures of shape:
  - ▶ All the quantiles together
  - ▶ Skew (beyond the scope of the class)
  - ▶ Kurtosis (beyond the scope of the class)

# Numerical summaries

## ▶ Numerical summary (statistic)

- ▶ A number or list of numbers calculated using the data (and only the data).
- ▶ Numerical summaries highlight important features of the data (shape, center, spread, outliers).

## ▶ Examples:

- ▶ Measures of center:
  - ▶ Arithmetic mean
  - ▶ Median
  - ▶ Mode
- ▶ Measures of spread:
  - ▶ Sample variance
  - ▶ Sample standard deviation
  - ▶ Range
  - ▶ IQR
- ▶ Measures of shape:
  - ▶ All the quantiles together
  - ▶ Skew (beyond the scope of the class)
  - ▶ Kurtosis (beyond the scope of the class)

# Numerical summaries

## ▶ Numerical summary (statistic)

- ▶ A number or list of numbers calculated using the data (and only the data).
- ▶ Numerical summaries highlight important features of the data (shape, center, spread, outliers).

## ▶ Examples:

- ▶ Measures of center:
  - ▶ Arithmetic mean
  - ▶ Median
  - ▶ Mode
- ▶ Measures of spread:
  - ▶ Sample variance
  - ▶ Sample standard deviation
  - ▶ Range
  - ▶ IQR
- ▶ Measures of shape:
  - ▶ All the quantiles together
  - ▶ Skew (beyond the scope of the class)
  - ▶ Kurtosis (beyond the scope of the class)

# Numerical summaries

## ▶ Numerical summary (statistic)

- ▶ A number or list of numbers calculated using the data (and only the data).
- ▶ Numerical summaries highlight important features of the data (shape, center, spread, outliers).

## ▶ Examples:

- ▶ Measures of center:
  - ▶ Arithmetic mean
  - ▶ Median
  - ▶ Mode
- ▶ Measures of spread:
  - ▶ Sample variance
  - ▶ Sample standard deviation
  - ▶ Range
  - ▶ IQR
- ▶ Measures of shape:
  - ▶ All the quantiles together
  - ▶ Skew (beyond the scope of the class)
  - ▶ Kurtosis (beyond the scope of the class)

# Numerical summaries

## ▶ Numerical summary (statistic)

- ▶ A number or list of numbers calculated using the data (and only the data).
- ▶ Numerical summaries highlight important features of the data (shape, center, spread, outliers).

## ▶ Examples:

- ▶ Measures of center:
  - ▶ Arithmetic mean
  - ▶ Median
  - ▶ Mode
- ▶ Measures of spread:
  - ▶ Sample variance
  - ▶ Sample standard deviation
  - ▶ Range
  - ▶ IQR
- ▶ Measures of shape:
  - ▶ All the quantiles together
  - ▶ Skew (beyond the scope of the class)
  - ▶ Kurtosis (beyond the scope of the class)

# Measures of center

$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$
0	1	1	2	3	5

- ▶ Arithmetic mean:
  - ▶  $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$
  - ▶ Here,  $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$
- ▶ Median:  $Q(0.5)$ .
  - ▶ A shortcut to calculating  $Q(0.5)$  is:
    - ▶  $Q(0.5) = x_{[n/2]}$  if  $n$  is odd
    - ▶  $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$  if  $n$  is even.
  - ▶ Here,  $Q(0.5) = (1 + 2)/2 = 1.5$
- ▶ Mode (of a discrete or categorical dataset)
  - ▶ the most frequently-occurring value
  - ▶ Here, mode = 1.

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

# Measures of center

$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$
0	1	1	2	3	5

- ▶ Arithmetic mean:

- ▶  $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$

- ▶ Here,  $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$

- ▶ Median:  $Q(0.5)$ .

- ▶ A shortcut to calculating  $Q(0.5)$  is:

- ▶  $Q(0.5) = x_{[n/2]}$  if  $n$  is odd

- ▶  $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$  if  $n$  is even.

- ▶ Here,  $Q(0.5) = (1 + 2)/2 = 1.5$

- ▶ Mode (of a discrete or categorical dataset)

- ▶ the most frequently-occurring value

- ▶ Here, mode = 1.

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

# Measures of center

$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$
0	1	1	2	3	5

- ▶ Arithmetic mean:

- ▶  $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$

- ▶ Here,  $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$

- ▶ Median:  $Q(0.5)$ .

- ▶ A shortcut to calculating  $Q(0.5)$  is:

- ▶  $Q(0.5) = x_{[n/2]}$  if  $n$  is odd

- ▶  $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$  if  $n$  is even.

- ▶ Here,  $Q(0.5) = (1 + 2)/2 = 1.5$

- ▶ Mode (of a discrete or categorical dataset)

- ▶ the most frequently-occurring value

- ▶ Here, mode = 1.

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

# Measures of center

$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$
0	1	1	2	3	5

► Arithmetic mean:

►  $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$

► Here,  $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$

► Median:  $Q(0.5)$ .

► A shortcut to calculating  $Q(0.5)$  is:

►  $Q(0.5) = x_{[n/2]}$  if  $n$  is odd

►  $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$  if  $n$  is even.

► Here,  $Q(0.5) = (1 + 2)/2 = 1.5$

► Mode (of a discrete or categorical dataset)

► the most frequently-occurring value

► Here, mode = 1.

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

# Measures of center

$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$
0	1	1	2	3	5

► Arithmetic mean:

►  $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$

► Here,  $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$

► Median:  $Q(0.5)$ .

► A shortcut to calculating  $Q(0.5)$  is:

►  $Q(0.5) = x_{[n/2]}$  if  $n$  is odd

►  $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$  if  $n$  is even.

► Here,  $Q(0.5) = (1 + 2)/2 = 1.5$

► Mode (of a discrete or categorical dataset)

► the most frequently-occurring value

► Here, mode = 1.

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

# Measures of center

$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$
0	1	1	2	3	5

- ▶ Arithmetic mean:
  - ▶  $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$
  - ▶ Here,  $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$
- ▶ Median:  $Q(0.5)$ .
  - ▶ A shortcut to calculating  $Q(0.5)$  is:
    - ▶  $Q(0.5) = x_{[n/2]}$  if  $n$  is odd
    - ▶  $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$  if  $n$  is even.
  - ▶ Here,  $Q(0.5) = (1 + 2)/2 = 1.5$
- ▶ Mode (of a discrete or categorical dataset)
  - ▶ the most frequently-occurring value
  - ▶ Here, mode = 1.

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

# Measures of center

$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$
0	1	1	2	3	5

- ▶ Arithmetic mean:
  - ▶  $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$
  - ▶ Here,  $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$
- ▶ Median:  $Q(0.5)$ .
  - ▶ A shortcut to calculating  $Q(0.5)$  is:
    - ▶  $Q(0.5) = x_{\lceil n/2 \rceil}$  if  $n$  is odd
    - ▶  $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$  if  $n$  is even.
  - ▶ Here,  $Q(0.5) = (1 + 2)/2 = 1.5$
- ▶ Mode (of a discrete or categorical dataset)
  - ▶ the most frequently-occurring value
  - ▶ Here, mode = 1.

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

# Measures of center

$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$
0	1	1	2	3	5

- ▶ Arithmetic mean:
  - ▶  $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$
  - ▶ Here,  $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$
- ▶ Median:  $Q(0.5)$ .
  - ▶ A shortcut to calculating  $Q(0.5)$  is:
    - ▶  $Q(0.5) = x_{\lceil n/2 \rceil}$  if  $n$  is odd
    - ▶  $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$  if  $n$  is even.
  - ▶ Here,  $Q(0.5) = (1 + 2)/2 = 1.5$
- ▶ Mode (of a discrete or categorical dataset)
  - ▶ the most frequently-occurring value
  - ▶ Here, mode = 1.

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

# Measures of center

$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$
0	1	1	2	3	5

- ▶ Arithmetic mean:
  - ▶  $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$
  - ▶ Here,  $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$
- ▶ Median:  $Q(0.5)$ .
  - ▶ A shortcut to calculating  $Q(0.5)$  is:
    - ▶  $Q(0.5) = x_{\lceil n/2 \rceil}$  if  $n$  is odd
    - ▶  $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$  if  $n$  is even.
  - ▶ Here,  $Q(0.5) = (1 + 2)/2 = 1.5$
- ▶ Mode (of a discrete or categorical dataset)
  - ▶ the most frequently-occurring value
  - ▶ Here, mode = 1.

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$
0	1	1	2	3	5

- ▶ Arithmetic mean:
  - ▶  $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$
  - ▶ Here,  $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$
- ▶ Median:  $Q(0.5)$ .
  - ▶ A shortcut to calculating  $Q(0.5)$  is:
    - ▶  $Q(0.5) = x_{\lceil n/2 \rceil}$  if  $n$  is odd
    - ▶  $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$  if  $n$  is even.
  - ▶ Here,  $Q(0.5) = (1 + 2)/2 = 1.5$
- ▶ Mode (of a discrete or categorical dataset)
  - ▶ the most frequently-occurring value
  - ▶ Here, mode = 1.

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

# Measures of center

$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$
0	1	1	2	3	5

- ▶ Arithmetic mean:
  - ▶  $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$
  - ▶ Here,  $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$
- ▶ Median:  $Q(0.5)$ .
  - ▶ A shortcut to calculating  $Q(0.5)$  is:
    - ▶  $Q(0.5) = x_{\lceil n/2 \rceil}$  if  $n$  is odd
    - ▶  $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$  if  $n$  is even.
  - ▶ Here,  $Q(0.5) = (1 + 2)/2 = 1.5$
- ▶ Mode (of a discrete or categorical dataset)
  - ▶ the most frequently-occurring value
  - ▶ Here, mode = 1.

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

# Measures of center

$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$
0	1	1	2	3	5

- ▶ Arithmetic mean:
  - ▶  $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$
  - ▶ Here,  $\bar{x} = \frac{1}{6}(0 + 1 + 1 + 2 + 3 + 5) = 2$
- ▶ Median:  $Q(0.5)$ .
  - ▶ A shortcut to calculating  $Q(0.5)$  is:
    - ▶  $Q(0.5) = x_{\lceil n/2 \rceil}$  if  $n$  is odd
    - ▶  $Q(0.5) = (x_{n/2} + x_{n/2+1})/2$  if  $n$  is even.
  - ▶ Here,  $Q(0.5) = (1 + 2)/2 = 1.5$
- ▶ Mode (of a discrete or categorical dataset)
  - ▶ the most frequently-occurring value
  - ▶ Here, mode = 1.

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

# Measures of spread

	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$
$x_i$	0	1	1	2	3	5
$\frac{i-.5}{n}$	.083	0.25	0.417	0.583	0.75	0.917

## ► Sample variance

- $s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$

- Here,  $s^2 = \frac{1}{6-1} [(0-2)^2 + (1-2)^2 + (1-2)^2 + (2-2)^2 + (3-2)^2 + (5-2)^2] = 3.2$

## ► Sample standard deviation

- $s = \sqrt{s^2} = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2}$

- Here,  $s = \sqrt{3.2} = 1.7889$

## ► Range

- Range = Maximum - Minimum

- Here, Range =  $5 - 0 = 5$

## ► Interquartile range

- $IQR = Q(0.75) - Q(0.25)$

- Here,  $IQR = 3 - 1 = 2$ .

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

# Your turn: sensitivity to outliers

Compare:

	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$
$x_i$	0	1	1	2	3	5
$\frac{i-.5}{n}$	.083	0.25	0.417	0.583	0.75	0.917

to:

	$y_1$	$y_2$	$y_3$	$y_4$	$y_5$	$y_6$
$x_i$	0	1	1	2	3	817263489
$\frac{i-.5}{n}$	.083	0.25	0.417	0.583	0.75	0.917

which measures of center and spread differ drastically between the  $x_i$ 's and the  $y_i$ 's? Which ones are about the same?

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

# Answers: sensitivity to outliers

Data	$x_i$	$y_i$
Mean	2	$1.3621 \times 10^8$
Median	1.5	1.5
Mode	1	1
Sample Variance	3.2	$1.1132 \times 10^{17}$
Sample Std. Dev.	1.7889	$3.3365 \times 10^8$
Range	5	$8.1726 \times 10^8$
IQR	2	2

# Sensitivity of numerical summaries

- ▶ Numerical summaries sensitive to outliers *and* skewness:
  - ▶ Mean
  - ▶ Sample variance
  - ▶ Sample standard deviation
  - ▶ Range
- ▶ Less sensitive numerical summaries:
  - ▶ Median
  - ▶ Mode
  - ▶ IQR

# Outline

Descriptive  
Statistics: Part  
2/2 (Ch 3)

Will Landau

Boxplots

Boxplots

Quantile-Quantile  
(QQ) Plots

Quantile-Quantile (QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Theoretical Quantile-Quantile Plots

Numerical  
Summaries

Numerical Summaries

Parameters

Parameters

# Statistics and parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.
  - ▶ Population mean (“true” mean):
    - ▶  $\mu = \frac{1}{N} \sum_{i=1}^N x_i$  if  $N$  the *finite* population size.
    - ▶  $\bar{x} \approx \mu$ .
  - ▶ Population variance (“true” variance):
    - ▶  $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$  if  $N$  the *finite* population size.
    - ▶  $s^2 \approx \sigma^2$ .
  - ▶ Population standard deviation (“true” standard deviation):
    - ▶  $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$  if  $N$  is the *finite* population size.
    - ▶  $s \approx \sigma$ .

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.

- ▶ Population mean ("true" mean):

- ▶  $\mu = \frac{1}{N} \sum_{i=1}^N x_i$  if  $N$  the *finite* population size.
  - ▶  $\bar{x} \approx \mu$ .

- ▶ Population variance ("true" variance):

- ▶  $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$  if  $N$  the *finite* population size.
  - ▶  $s^2 \approx \sigma^2$ .

- ▶ Population standard deviation ("true" standard deviation):

- ▶  $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$  if  $N$  is the *finite* population size.
  - ▶  $s \approx \sigma$ .

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.

- ▶ Population mean (“true” mean):

- ▶  $\mu = \frac{1}{N} \sum_{i=1}^N x_i$  if  $N$  the *finite* population size.
- ▶  $\bar{x} \approx \mu$ .

- ▶ Population variance (“true” variance):

- ▶  $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$  if  $N$  the *finite* population size.
- ▶  $s^2 \approx \sigma^2$ .

- ▶ Population standard deviation (“true” standard deviation):

- ▶  $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$  if  $N$  is the *finite* population size.
- ▶  $s \approx \sigma$ .

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.
  - ▶ Population mean (“true” mean):
    - ▶  $\mu = \frac{1}{N} \sum_{i=1}^N x_i$  if  $N$  the *finite* population size.
    - ▶  $\bar{x} \approx \mu$ .
  - ▶ Population variance (“true” variance):
    - ▶  $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$  if  $N$  the *finite* population size.
    - ▶  $s^2 \approx \sigma^2$ .
  - ▶ Population standard deviation (“true” standard deviation):
    - ▶  $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$  if  $N$  is the *finite* population size.
    - ▶  $s \approx \sigma$ .

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.
  - ▶ Population mean (“true” mean):
    - ▶  $\mu = \frac{1}{N} \sum_{i=1}^N x_i$  if  $N$  the *finite* population size.
    - ▶  $\bar{x} \approx \mu$ .
  - ▶ Population variance (“true” variance):
    - ▶  $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$  if  $N$  the *finite* population size.
    - ▶  $s^2 \approx \sigma^2$ .
  - ▶ Population standard deviation (“true” standard deviation):
    - ▶  $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$  if  $N$  is the *finite* population size.
    - ▶  $s \approx \sigma$ .

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.
  - ▶ Population mean (“true” mean):
    - ▶  $\mu = \frac{1}{N} \sum_{i=1}^N x_i$  if  $N$  the *finite* population size.
    - ▶  $\bar{x} \approx \mu$ .
  - ▶ Population variance (“true” variance):
    - ▶  $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$  if  $N$  the *finite* population size.
    - ▶  $s^2 \approx \sigma^2$ .
  - ▶ Population standard deviation (“true” standard deviation):
    - ▶  $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$  if  $N$  is the *finite* population size.
    - ▶  $s \approx \sigma$ .

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.
  - ▶ Population mean (“true” mean):
    - ▶  $\mu = \frac{1}{N} \sum_{i=1}^N x_i$  if  $N$  the *finite* population size.
    - ▶  $\bar{x} \approx \mu$ .
  - ▶ Population variance (“true” variance):
    - ▶  $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$  if  $N$  the *finite* population size.
    - ▶  $s^2 \approx \sigma^2$ .
  - ▶ Population standard deviation (“true” standard deviation):
    - ▶  $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$  if  $N$  is the *finite* population size.
    - ▶  $s \approx \sigma$ .

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.
  - ▶ Population mean (“true” mean):
    - ▶  $\mu = \frac{1}{N} \sum_{i=1}^N x_i$  if  $N$  the *finite* population size.
    - ▶  $\bar{x} \approx \mu$ .
  - ▶ Population variance (“true” variance):
    - ▶  $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$  if  $N$  the *finite* population size.
    - ▶  $s^2 \approx \sigma^2$ .
  - ▶ Population standard deviation (“true” standard deviation):
    - ▶  $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$  if  $N$  is the *finite* population size.
    - ▶  $s \approx \sigma$ .

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.
  - ▶ Population mean (“true” mean):
    - ▶  $\mu = \frac{1}{N} \sum_{i=1}^N x_i$  if  $N$  the *finite* population size.
    - ▶  $\bar{x} \approx \mu$ .
  - ▶ Population variance (“true” variance):
    - ▶  $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$  if  $N$  the *finite* population size.
    - ▶  $s^2 \approx \sigma^2$ .
  - ▶ Population standard deviation (“true” standard deviation):
    - ▶  $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$  if  $N$  is the *finite* population size.
    - ▶  $s \approx \sigma$ .

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.
  - ▶ Population mean (“true” mean):
    - ▶  $\mu = \frac{1}{N} \sum_{i=1}^N x_i$  if  $N$  the *finite* population size.
    - ▶  $\bar{x} \approx \mu$ .
  - ▶ Population variance (“true” variance):
    - ▶  $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$  if  $N$  the *finite* population size.
    - ▶  $s^2 \approx \sigma^2$ .
  - ▶ Population standard deviation (“true” standard deviation):
    - ▶  $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$  if  $N$  is the *finite* population size.
    - ▶  $s \approx \sigma$ .

Boxplots

Quantile-Quantile  
(QQ) Plots

Theoretical  
Quantile-Quantile  
Plots

Numerical  
Summaries

Parameters

- ▶ **Statistic:** numerical summary of data on the *sample*
- ▶ **Parameter:** numerical summary of a theoretical distribution or data on an entire *population*.
  - ▶ Population mean (“true” mean):
    - ▶  $\mu = \frac{1}{N} \sum_{i=1}^N x_i$  if  $N$  the *finite* population size.
    - ▶  $\bar{x} \approx \mu$ .
  - ▶ Population variance (“true” variance):
    - ▶  $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$  if  $N$  the *finite* population size.
    - ▶  $s^2 \approx \sigma^2$ .
  - ▶ Population standard deviation (“true” standard deviation):
    - ▶  $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$  if  $N$  is the *finite* population size.
    - ▶  $s \approx \sigma$ .